



JUARA: Jurnal Olahraga

E-ISSN 2655-1896 ISSN 2443-1117
<https://doi.org/10.33222/juara.v5i1.1000>



Age 6-9 Years: Nutritional Status on Children's Physical Fitness

Sapika^{1*}, Rasyidah Jalil^{2*}, Wahyuni Ulpi^{3*}, Nurwahidin Hakim⁴

¹Muhammadiyah University of Palopo, Indonesia

²Muhammadiyah University of Palopo, Indonesia

³Muhammadiyah University of Palopo, Indonesia

⁴Muhammadiyah University of Palopo, Indonesia

*e-mail:

sapikaa@student.umpalopo.ac.id

rasyidah@umpalopo.ac.id

wahyuniulpi@umpalopo.ac.id

nurwahidinhakin@umpalopo.ac.id

Article Info

Article History:

Received (month) (year)

Approved (month) (year)

Published (month) (year)

Keywords:

Nutritional Status,

Physical Fitness

Abstract

The purpose of this study was to determine whether there is a relationship between children's nutritional status and physical fitness. This research is a quantitative research with a correlational research design whose goal is linking two or more variables. The instrument used to measure the nutritional status of children is the maximum physical fitness index (BMI) and the Indonesian physical fitness test (TKJI). The results of the data were analyzed using the SPSS application. The results of the descriptive analysis of the nutrition variable had the lowest score of 13.3 and the highest score of 19.2 with an average value of 18.8 while the TKJI variable had the lowest score of 13 and the highest score of 19 with an average value of 1.9. From the research data and analysis conducted, it can be concluded that the significant value is $0.002 < 0.05$. Thus, it means that there is a relationship between nutritional status and physical fitness with the meaning of H^1 received.

© 2019 STKIP Muhammadiyah Kuningan

Under the license CC BY-SA 4.0

✉Correspondence address:

E-mail :

INTRODUCTION

Physical fitness is one of the things that is expected by every individual because with physical fitness a person will perform optimally and confidently. (Wahyuni Ulpi, Nurwahidin Hakim, 2022). Physical fitness is a person's ability to complete a certain task or physical activity without getting tired. Many children at school are physically weak and tired when participating in activities. Students must be physically fit to participate in learning activities and ensure they are not sick or tired while studying. Physical fitness is a need that must be fulfilled by every human being to carry out daily activities as well as possible, in other words that physical fitness is the body's ability to carry out activities without experiencing excessive fatigue. The term physical fitness has a meaning that is no different from the physical aspects of total fitness. or what is known as physical fitness (Forms et al., 2020). Someone who has good physical fitness can be interpreted as having enough ability to do his job efficiently without causing significant fatigue, so that he still has energy left to fill his spare time and other sudden tasks. It can also be said that a good level of physical fitness gives a person the ability to lead a productive life and be able to adapt to many loads. Parents and schools are responsible for ensuring that students have a balanced diet. (Idham et al., 2022). Physical fitness is the goal of one of the implementation of sports and health physical education subjects, physical fitness basically functions as an integrated component of the wider education system, focusing on developing moral behavior, mental and emotional stability, and social and emotional abilities through activities physique. (Judge & Hidayat, 2020). Physical fitness is the basis for carrying out an activity as well as teaching so it really needs to be improved, especially in early childhood aged 6-9 years (Hakim et al., 2022). In carrying out activities, the higher the physical fitness, the easier it is to carry out activities (Ade Evriansyah Lubis, 2020). Vice versa, the worse the physical fitness, the more difficult it is to do activities. (Syaleh et al., 2019). Physical education is the contribution of part of the education program in general, especially through movement experiences to ensure the growth and development of children. Physical education is an integral part of education through movement experiences that encourage

physical abilities, motor skills, cognitive development, social-emotional and spiritual development. An effective physical education process will accelerate physical education goals that have been designed such as physical development, movement development, movement skills, cognitive and affective development, social development and emotional development. The development of motion is one of the most important parts of the goal in implementing physical education (Nugraha, 2015).

Nutritional status

Food intake is all forms of food that are digested by the digestive system. With proper food intake, by not consuming the wrong food, such as consuming very large amounts of food, high in fat, excessive carbohydrates, and low in fiber, but not balanced with balanced energy expenditure, this is one of the causes of overweight or overweight cases. Body (Hita et al., 2020). In general, nutritional status is very important to determine health and balance between physical and mental growth. Factors related to physical fitness in general can be broken down into various categories, including: Heredity [1], Regular exercise [2], Gender [3], Age [4], Diet [5], Rest [6], Activity [7], and many other factors (Muliani, 2020). The child's weight and height are used to determine nutritional status, namely the balance of nutrients needed by the child's body. (1 & , Dary 2, 2022). Nutritional status is a global problem, inadequate nutrition can cause abnormalities in the growth and development of children (Dewi et al., 2022). Nutritional status is expressed as the quality of future human resources (Rahadiyanti et al., 2022). Nutritional status has four groups including poor nutrition, undernutrition, good nutrition, and excess nutrition which is used to describe the state of the body due to the use of nutrients and food consumption (Adi et al., 2019). Nutrition for early childhood is a key factor in encouraging the growth and development of children in the future (Widiastiwi et al., 2022).

Children aged 6 to 9 years are still included in the PAUD category based on the Constitution (Children & Years, 2019). School age is an age that requires more concern in terms of nutrition and health. At this age, children are able to consume the food and drink choices they want. (Hanim et al., 2022). School-age children are involved in a variety

of mental and physical activities, including playing, learning, and exercising. The nutrients offered will help improve the body's health so that the immune system can mature properly and prevent disease. (1 & , Dary 2, 2022). Consuming foods that offer nutrients is also needed by the body to maintain physical health. However, if a person's level of physical fitness is low, it is likely that they are eating foods that lack the nutrients their bodies need. (Patten & Newhart, 2018). The process of child growth and development is greatly influenced by excess nutrition, especially in the field of emotional psychology. This increases the risk of degenerative and/or metabolic disorders and the possibility of becoming obese as an adult. (Safitri et al., 2022). The main nutritional problems in Indonesia consist of basic nutritional problems, namely Protein Energy Deficiency (KEP), Vitamin A Deficiency (KVA), Disorders Due to Iodine Deficiency (IDD), and Iron Nutrition Anemia (AGB), in addition to excess nutrition (obesity). Indonesia is currently experiencing 2 nutritional problems at the same time or better known as the multiple nutrition problem. The handling of nutrition problems is closely related to a nation's strategy in creating healthy, intelligent and productive human resources. Efforts to improve quality human resources begin with handling the growth of children as part of a family with good nutrition and care (Pawitra et al., 2019).

Malnutrition, especially in the nutritionally vulnerable age group, can cause failure to thrive and increase mortality and morbidity (Novela & Kartika, 2019). Malnutrition can increase the risk of infectious diseases, slow down growth and development and reduce the level of intelligence, both undernutrition and over nutrition can disrupt the process of child growth and development (Novianti & Utami, 2021). If left untreated, malnutrition can lead to death, disability, and stunted physical growth, and ultimately disrupt the country's socio-economic development. (Riaz et al., 2021)

Direct and indirect factors greatly affect children, direct factors include: lack of culinary intake, as well as infectious diseases, while indirect factors are parents' income, food availability at the place of residence, diet, environmental sanitation, health services, parents' occupation, and knowledge that ultimately susceptible to infectious diseases.

(Rahmasari et al., 2022). Furthermore, birth spacing that is too close also affects the nutritional status of children. (Herlambang et al., 2021). Factors that work with physical fitness can generally be broken down into various categories, including Heredity, Regular exercise, Gender, Age, Diet, Rest, physical activity, and many other factors (1 & , Dary 2, 2022).

Improvements in nutrition are needed starting from pregnancy, infants and toddlers, preschoolers, elementary school-age children, adolescents and adults, to old age. Elementary school children are a strategic target in improving community nutrition because during childhood the functions of the brain organs begin to form steadily so that the development of intelligence is quite rapid. Elementary School Children (SD) are children aged 6-12 years (Seprianty et al., 2015) Nutritional problems cannot be handled through short-term and sectoral policies and programs, let alone only from a food perspective. Nutritional problems must be addressed immediately through the implementation of appropriate nutrition policies as a whole. (Ernawati et al., 2019). Stunted nutrition in children is not only a health sector problem but also a multisector problem. (Adam & Medong, 2022).

One of the factors that affect a person's nutrition is a lack of knowledge about nutrition. This reduced knowledge will also reduce a person's ability to apply nutritional information in everyday life. One way to increase someone's knowledge is by providing nutrition education as early as possible. Nutrition education can be provided through counseling, giving posters, leaflets or booklets to school children (Rahadiyanti et al., 2022). Nutrition education for anemic children in elementary schools is given with the hope that the child's nutritional knowledge and the child's eating pattern will change so that the child's food intake, especially iron intake, will be better. With better iron intake, the child's hemoglobin level will increase (Novian, 2013)

METHODS

This type of research uses quantitative research, where this research applies methods for direct data collection (Hardiansyah & Syampurma, 2017). Quantitative research uses SPSS data management. Quantitative research has the goal of generalizing research findings so that they can be used to predict the same situation in other populations. Quantitative

research is also used to explain the causal relationship between the variables studied. Quantitative research begins with theories and hypotheses. Researchers use manipulation techniques and control variables through formal instruments to see causality interactions (Abdullah, 2015)

The sample is a small part of the population and the sampling or data source is done purposively and snowball. The subjects of this study consisted of 10 children, including 5 children and 5 girls. This research was conducted in the sub-district of Bua, Luwu Regency. In this study, the nutritional status of children was analyzed for physical fitness. using Body Mass Index (BMI) measurements while the data collection mechanism on the level of physical fitness of students uses the Indonesian Physical Fitness Test (TKJI) for the age range of 6-9 years, the test items include running 30 meters, pull ups (hanging elbows), vertical jump, (jump straight), sit up (get up sitting) and run 600 meters, the instrument used is a height measuring meter, weight scales, stopwatch., what will happen in the research will be analyzed using the relationship technique and also using the correlation analysis technique, in addition to using the percentage analysis technique to see illustrations of data in general using the SPSS application (Saparia et al., 2022). Correlation is a reciprocal or causal relationship in other words that correlation is to see the extent to which a child's nutritional status is related to physical fitness, so that the data analysis process requires critical thinking because it will produce real data and can be used as a reference for further research. Data analysis descriptive Analysis Finding the average or Mean of Variable X and Variable Y is the process of processing data with the aim of finding useful information that can be used as a basis for making decisions to solve a problem. with the following formula :

$$Mx/\bar{X} = \frac{\sum \bar{X}}{N}$$

$$My/\bar{Y} = \frac{\sum \bar{Y}}{N}$$

No	Total Value	Classification
1	22-25	Excellent

2	18 – 21	Good
3	14 – 17	Fairly Good
4	10 – 13	Bad
5	5 – 9	Very Bad

Source : (Mashuri & Pasaribu, 2019)

FINDINGS AND DISCUSSION

The results of the research are then analyzed using descriptive analysis to discover an overview of students' abilities on the two test items carried out, which are the body mass index test and the physical fitness test (TKJI). The research results can be seen in the following table:

Table 1. Analysis of Frequency Distribution of Physical Freshness Test Results for Children Aged 6-9 Years.

Category	Frequency	Percentage
Excellent	0	0%
Good	2	20%
Fairly Good	8	80%
Bad	0	0%
Very Bad	0	0%
Total	10	100%

The results of the data from each item of physical fitness test after categorization which was then carried out by percentage analysis found that out of 10 samples most were in the "Fairly Good" category, namely 80% or as many as 8 children and the "good" category was only 20%.

Table 2. Frequency of Distribution Analysis of Children's Nutritional Status Measurement Results.

CATEGORY	FREQUENCY	PERCENTAGE
Malnutrition (thin)	0	0%
Good Nutrition (normal)	9	90%
Over nutrition (overweight)	1	10%
Obesity (>+2SD)	0	0%
TOTAL	10	100%

The results of measuring the nutritional status of children were then analyzed using the body mass index (BMI) formula, the results of 10 children were mostly in the "normal" nutrition category, namely 9

people or 90% of the total sample, while the rest were in the overweight category. or over nutrition, namely 1 student or 10%.

relations between nutritional status and physical fitness because the significance value is <0.05, meaning that H1 is accepted.

DESCRIPTIVE ANALYSIS

Statistics

		nutrition	TKJI
N	Valid	10	10
	missing	0	0
Means		16,160	15.50
Median		15,800	15.00
std. Deviation		1.9062	1,958
Range		5,9	6
Minimum		13,3	13
Maximum		19,2	19
sum		161.6	155

Based on descriptive analysis with a sample of 10 consisting of 5 boys and 5 girls In the table above it can be seen that the nutrition variable has the lowest score of 13.3 and the highest score of 19.2 with an average score of 15.8 and a standard deviation (level of data distribution) 1.90. The TKJI variable has the lowest score of 13 and the highest score of 19, with an average value of 15 with a standard deviation of 1.9.

REGRESSION TEST

Model	Unstandardized Coefficients		Standardized Coefficients	t	Sig.
	B	std. Error			
1 (Constant)	1,566	3,208		,488	,639
nutrition	,862	,197	,840	4,371	,002

In the table coefficients which in column B the constant (a) is 1.566, while the trust value (b) is 0.862, thus the regression equation can be interpreted: $Y=a + bX$ or $1.566 + 0.862$ with a significant value of 0.002. So it can be concluded that there is a

DISCUSSION

1. Categories of nutritional status in children aged 6-9 years.

The description of the Nutritional Status of Children shows that the highest percentage of Nutritional Status is in the (Normal) category of 9 people 90% while the rest are in the Over Weight or Overnutrition category, which only 1 person or 10% School-age children need more concern in terms of Nutrition and Health, at the age of 6-9 years children have been able to consume the food and drink choices they want. This is supported by research(Ferida, 2012)said that nutritional status affects the physical fitness of children.

Nutritional status in the "Normal" category indicates that most of the respondents have sufficient intake of nutrients that enter the child's body. This is in line with using research at SD N Batur 2 and SDN Tuguran Nogotirto using a sample of 100 children with the (Normal) category of 73%, this shows that the balance of food intake greatly influences the growth of children, especially elementary school children, this growth will greatly affect to enter the next stage(Haryani et al., 2020)

Children classified as Nutrition (Normal) at SDN Sumber Sekar 1 Kec. Dau City of Malang gets a percentage of 60% and 12 children included in the Nutrition criteria (thin) get a percentage of 34%, with a sampling technique that is purposive sampling(Wahyudi et al., 2017)

This study explains that the results of examining the nutritional status of students at SDN 1 Piton used the Body Max Index (BMI) where 60 respondents, 27 respondents with a percentage of 45% were under nutrition (fat) and 33 respondents were undernourished (thin) with a percentage of 55%.(Zahra et al., 2020).

2. Physical fitness test category.

The description of the child's Physical Fitness test shows that the percentage of the physical fitness test is in the category "Very good" there are no students in the achievement or 0%, the category "Good" is 20%, the category "Moderate" is 80% or as many as 8 people, the category "Less" 0%, category "Not very much" 0%.

Achievement of physical fitness results at SDN Pahrubuh 1 and MI Mambaul Hikam, Kediri District, Semen District, Kediri Regency, the highest score was in the "Medium" category of 11 students, the "Less" category was achieved by 9 students, the "Good" category was achieved by 2 students in the "Less" category. once" was achieved by 2 students, and in the "very good" category there were no students who reached that category. (Wirnantika et al., 2017)

Physical fitness in children is not only determined by nutritional status but in physical activity according to measurements of the level of physical fitness providing daily activities and can affect the quality of Physical Fitness. Physical fitness of students can be predicted through physical activity measured using a physical fitness test. The better the child's physical activity, the higher the child's physical fitness.

3. Relationship between nutritional status of children aged 6-9 years to physical fitness

The SPSS test has been carried out to see the relationship between children's nutritional status and physical fitness the research data and analysis carried out concluded that the significant value was $0.002 < 0.05$ so that it can be said that there is a relationship between nutritional status and physical fitness, meaning that H1 is accepted and H0 is rejected, which means there is a relationship between nutritional status and physical fitness of children aged 6-9 years .

CONCLUSION

The results of the research and discussion show that in general 90% of children are seen from the nutritional status in the (Fairly Good) category and 80% in the physical fitness test are in the (Good) category. In addition, there is a significant relationship between nutritional status and physical fitness. Physical fitness status can be supported by balanced nutritional intake. It has been proven that nutritional status works the same as physical fitness. Balanced nutritional status has an impact on physical fitness, eating patterns and improper nutritional intake will cause nutritional problems in children. It was found that factors influencing the attainment of nutritional status included parents' income,

parents' knowledge of nutrition, individual illnesses, and physical growth.

ACKNOWLEDGMENTS

Thank you to those who have helped, especially the University of Muhammadiyah Palopo for helping and facilitating us in the research process to the end and to the JUARA: Jurnal Olahraga.

REFERENCES

- 1, MSM, & , Dary 2, DCD 3. (2022). Muhammadiyah Journal of Nursing. J Journal of Nursing Muhammadiyah Address, 7(1), 3–6.
- Abdullah, PM (2015). Living in the world that is fit for habitation : CCI's ecumenical and religious relationships. In Aswaja Pressindo.
- Adam, G., & Medong, A. (2022). Nutritional Problems, Stunting and Their Impact on Early Childhood Growth and Development. Lonto Leok Journal, 4(1), 1–10.
- Ade Evriansyah Lubis, AN (2020). Journal of Teaching Physical Education in Elementary Schools. Journal of Teaching Physical Education in Elementary Schools, 4(77), 73–80.
- Adi, M., University, P., & Malang, N. (2019). The Relationship between Nutritional Status and Physical Activity with the Level of Physical Fitness in Junior High School Students. 1(3), 138–142.
- Children, P., & Years, U. (2019). 10.5281/Zenodo.3338697. 5(2), 11–17.
- Form, P., Squat, L., Against, J., Shooting, K., & Ball, S. (2020). MensSana Journal MensSana Journal. 5, 182–190.
- Dewi, TS, Widiastuti, S., & Argarini, D. (2022). Relationship between Parenting and Mother's Education with the Nutritional Status of Toddler-aged Children in the Gang Langgar Petogogan RW 03. Malahayati Nursing Journal, 4(3), 613–626. <https://doi.org/10.33024/mnj.v4i3.6037>

- Ernawati, A., Planning, B., Regional, P., Pati, K., Raya, J., Km, P.-K., & Tengah, P. 59163 J. (2019). Analysis of the Implementation of the Program for Overcoming Malnutrition in Toddlers at the Jakenan Health Center, Pati Regency. *Journal of R&D*, XV(1), 39–50.
- Ferida, PRA (2012). Correlation of Nutritional Status with the Physical Freshness Level of Elementary School Children. *Journal of Physical Education, Sport, Health and Recreations*, 1(1), 11–15.
- Fieny, T. et. a. (2016). *88 Care Journal* Vol. 4, No.3, Year 2016. 4(3), 88–96.
- Hakim, N., & Hidayat, R. (2020). The Relationship between Physical Fitness and the Ability to Dribble a Ball in Students of SDN 65 Pajalesang Palopo. *Journal of Sport and Physical Education*, 1(1), 59–66.
- Hakim, N., Ulpi, W., & Hasyim, FA (2022). Improving Children's Physical Fitness during a Pandemic through Windows Movie Maker-Based Gymnastics Activities. 6(4), 2677–2689. <https://doi.org/10.31004/obsession.v6i3.1954>
- Hanim, B., Ingelia, I., & Ariyani, D. (2022). Breakfast Habits with Nutritional Status of Elementary School Children. *Malakbi Midwifery Journal*, 3(1), 28. <https://doi.org/10.33490/b.v3i1.570>
- Hardiansyah, S., & Syampurma, H. (2017). Differences in Physical Freshness Levels Based on Nutritional Status. 7.
- Haryani, W., Setiyobroto, I., & Siregar, IHY (2020). The Influence of the Knowledge about Cariogenic Food Towards Dental Caries and Nutrition Status among 9-11 Years Old Children. *Journal of Dental Health*, 7(1), 40–45. <https://doi.org/10.31983/jkg.v7i1.5674>
- Herlambang, A., Wandini, R., & Setiawati. (2021). Factors Affecting the Nutritional Status of Toddlers in Indonesia. *JKM (Malayayati Midwifery Journal)*, 7(4), 673–680.
- Hita, IPAD, Ariestika, E., Billy Yacs, BTPW, & Pranata, D. (2020). Relationship of Nutritional Status to PMI Physical Activity Levels During the Covid-19 Quarantine Period. *MensSana Journal*, 5(2), 146–156. <https://doi.org/10.24036/menssana.050220.07>
- Idham, Z., Neldi, H., Komaini, A., Sin, TH, & Damrah, D. (2022). Effect of Physical Fitness, Nutritional Status, and Learning Motivation on PJOK Learning Outcomes. *Basicedu Journal*, 6(3), 4078–4089. <https://doi.org/10.31004/basicedu.v6i3.2196>
- Mashuri, H., & Pasaribu, AMN (2019). The Role of Rhythmic Gymnastics Against Physical Fitness for Elementary School Students. *SPORTIF Journal: Learning Research Journal*, 5(1), 89–97.
- Muliani, D. (2020). Relationship between Nutritional Status and Physical Fitness Level of Students of SMP Negeri 3 X Koto Singkarak, Solok Regency. 01(01), 1–6.
- Novela, V., & Kartika, L. (2019). Factors of Undernutrition Status in Preschool Children in the Work Area of the Guguk Panjang Health Center, Bukittinggi City. *Journal of Endurance*, 4(2), 359. <https://doi.org/10.22216/jen.v4i2.4021>
- Novian, A. (2013). *Andalas Public Health Journal*. Central Obesity And Total Blood Cholesterol Levels, 9(1), 37–43.
- Novianti, A., & Utami, TP (2021). Assessment of Nutritional Status and Balanced Nutrition Knowledge of School-Age Children as a Form of Activation of UKS Activities. *ABDIMAS: Journal of Community Service*, 4(1), 399–404. <https://doi.org/10.35568/abdimas.v4i1.908>
- Nugraha, B. (2015). Early Childhood Sports Physical Education. *Journal of Children's Education*, 4(1), 557–564. <https://doi.org/10.21831/jpa.v4i1.12344>

- Patten, ML, & Newhart, M. (2018). Sampling Clusters. *Understanding Research Methods*, 98–99. <https://doi.org/10.4324/9781315213033-32>
- Pawitra, PRA, Susilo, S., & Kusmawati, W. (2019). Development of a Textbook of Sports Nutrition for IKIP Budi Utomo Malang Students: Study of Critical Thinking Skills. *MensSana Journal*, 4(2), 175. <https://doi.org/10.24036/jm.v4i2.106>
- Rahadiyanti, A., Salma Dina, S., Putri, S., Tampubolon, O., Yesi Veicinlun, S., Mattarahmawati, SA, Jannah, SR, & Ningsih, S. (2022). Increasing Knowledge of Mothers and Teachers Regarding Balanced Nutrition for Children Aged 5-8 Years at Al-Hunafa School, Bandung City. *Ejournal2.Undip.Ac.Id*, 2022(1), 8–14.
- Rahmasari, Y., Anggraeni, S., Rahman, E., Studies, P., Society, K., & Society, FK (2022). FACTORS RELATED TO THE NUTRITIONAL STATUS OF TODDLERS AT THE CEMAKA PUTIH HEALTH CENTER, BANJARMASIN CITY IN 2022.
- Riaz, M., Azam, N., Mehmood, H., Asif, R., Khan, N., Ali, F., & Mughal, R. (2021). Nutritional status assessments smentoforphanagechildren *reninrawalpind i*. 71(6), 2139–2144.
- Safitri, DE, Fitri, ZE, & Lestari, WA (2022). Energy Consumption Density and Protein Intake Density Related to Nutritional Status of School-Age Children. *Muhammadiyah Journal of Nutrition and Food Science (MJNF)*, 2(2), 95. <https://doi.org/10.24853/mjnf.2.2.95-102>
- Saparia, A., Iskar, I., & Abduh, I. (2022). Analysis of the relationship between nutritional status and physical fitness level of adolescents aged 16-17 years. *Journal of Sports Science*, 4(2), 126. <https://doi.org/10.26418/jilo.v4i2.52540>
- Seprianty, V., Tjekyan, S., & Thaha, A. (2015). Nutritional Status of Class III Children at SDN 1 Sungaililin. *Journal of Medicine and Health*, 2(1), 129–134.
- Syaleh, M., Lubis, AE, Helmi, B., North, S., Education, S., & Health, J. (2019). The Scientific Journal of STOCK Bina Guna Medan LEARN TO SWIMMING FREE STYLE The Scientific Journal of STOK Bina Guna Medan. 7(4), 30–39.
- Wahyudi, DD, Yuliwar, R., & Maemunah, N. (2017). Differences in Nutritional Status of Elementary School Children Affected by Dental Caries and Not Dental Caries at Sumber Sekar 01 Public Elementary School, Dau District, Malang City. *Journal of Nursing News*, 2(1), 88–97.
- Wahyuni Ulpi, Nurwahidin Hakim, AK (2022). Overview of Physical Fitness for Early Childhood in. 6(1), 30–37. <https://doi.org/10.31004/obsession.v6i1.1197>
- Widiastiwi, Y., Zaidiah, A., & Ernawati, I. (2022). Monitoring the Nutritional Status of Early Childhood Education, Through the Utilization of Health and Information Technology. *Ikra-Ith Abdimas*, 5(2), 122–128.
- Wirmantika, I., Pratama, BA, & Hanief, YN (2017). Survey of Physical Fitness Levels of Grade IV Students at SDN Puhrubuh I and MI Mambaul Hikam in Kediri District for the 2016/2017 Academic Year. *Sportive*, 3(2), 240. https://doi.org/10.29407/js_unpgri.v3i2.11898
- Zahra, IM, Hidayati, S., & Mahirawatie, IC (2020). The Relationship between Nutritional Status and DMF-T in Students of SD Negeri 1 Piton, Punung District, Pacitan Regency. *Journal of Health Scales*, 11(2), 67–74. <https://doi.org/10.31964/jsk.v11i2.239>

